

INVADE-Publikationen

1. De Graauw J, Chonchol M, Poppert H, Etgen T, Sander D. Relationship between kidney function and risk of asymptomatic peripheral arterial disease in elderly subjects. *Nephrol Dial Transplant*. 2010 Jul 28
2. Kendrick J, Chonchol M, Gnahn H, Sander D. Higher systolic blood pressure is associated with progression of carotid intima-media thickness in patients with chronic kidney disease. *Kidney Int*. 2010 May; 77(9):794-800.
3. Sander K, Bickel H, Förstl H, Etgen T, Briesenick C, Poppert H, Sander D. Carotid- intima media thickness is independently associated with cognitive decline. The INVADE study. *Int J Geriatr Psychiatry*. 2010 Apr;25(4):389-94.
4. Ulm K, Huntgeburth U, Gnahn H, Briesenick C, Pürner K, Middeke M. Effect of an intensive nurse-managed medical care programme on ambulatory blood pressure in hypertensive patients. *Arch Cardiovasc Dis*. 2010 Mar;103(3):142-9
5. Etgen T, Sander D, Huntgeburth U, Poppert H, Förstl H, Bickel H. Physical activity and incident cognitive impairment in elderly persons: the INVADE study. *Arch Intern Med*. 2010 Jan 25; 170(2):186-93.
6. Etgen T, Sander D, Chonchol M, Briesenick C, Poppert H, Förstl H, Bickel H. Chronic kidney disease is associated with incident cognitive impairment in the elderly: the INVADE study. *Nephrol Dial Transplant*. 2009 Oct; 24(10):3144-50.
7. Schulze Horn C, Ilg R, Sander K, Bickel H, Briesenick C, Hemmer B, Poppert H, Sander D. High-sensitivity C-reactive protein at different stages of Atherosclerosis: results of the INVADE study. *J Neurol*. 2009 May; 256(5):783-91.
8. Chonchol M, Gnahn H, Sander D. Impact of subclinical atherosclerosis on incident chronic kidney disease in the elderly. *Nephrol Dial Transplant*. 2008 Aug; 23(8):2593-8.

9. Desbien AM, Chonchol M, Gnahn H, Sander D. Kidney Function and Progression of Carotid Intima-Media Thickness in a Community Study. *Am J Kidney Dis* 2008 Apr; 51(4):584-593.
10. Sander K, Bickel H, Schulze Horn C, Huntgeburth U, Poppert H, Sander D. Peripheral arterial disease: predictors and treatment intensity. Two-years of data from the population-based INVADE project. *Dtsch Med Wochenschr.* 2008 Mar; 133(10):455-9.
11. Schulze Horn C, Sander K, Ilg R, Bickel H, Briesenick C, Sander D. Modification of vascular risk factors by intervention on the primary care level. Results of the INVADE project. *Nervenarzt.* 2007 Dec; 78(12):1413-9.
12. Sander K, Horn CS, Briesenick C, Sander D; INVADE Study Group. High-sensitivity C-reactive protein is independently associated with early carotid artery progression in women but not in men: the INVADE Study. *Stroke.* 2007 Nov; 38(11):2881-6.
13. Sander K, Schulze Horn C, Briesenick C, Sander D. Significant reduction of vascular risk factors within the framework of the population-based intervention project INVADE--a 2-year follow-up study. *Dtsch Med Wochenschr.* 2006 Aug 25;131(34-35):1853-9.
14. Sander D, Schulze-Horn C, Bickel H, Gnahn H, Bartels E, Conrad B. Combine effects of hemoglobin A1c and C-reactive protein on the progression of subclinical carotid atherosclerosis: the INVADE study. *Stroke.* 2006 Feb;37(2):351-7.
15. Scherpinski U, Bickel H, Gnahn H, Förstl H, Conrad B, Sander D. Intervention project on cerebrovascular diseases and dementia in the Ebersberg district (INVADE): rationale and design. *Nervenarzt.* 2002 Dec; 73(12):1199-204.

Aktualisiert am 27.10.2010